

United States Patent Application

of

Dr. Klaus-Ulrich Weithmann

For

A Method for Detecting Active Ingredients of Medical Value

09876091.060801

chemical conversion of the substrate. Absent or reduced conversions of the marker substrate can also indicate the presence of said inhibitors. It is substantially more difficult to find inhibitors for the binding domain.

- 5 It is therefore the object of the present invention to find substances which reduce or essentially prevent binding of substrates to the binding domain of a protein. Said substance may act as inhibitor or may be a ligand.

- 10 The object is achieved by incubating a protein, the marker substrate and the substrate with a substance and by determining whether the marker substrate is converted by the protein.

The invention therefore relates to a method to determine whether a test substance is an inhibitor or a ligand of a protein, which method comprises

- 15 a) using a protein which contains at least one catalytic domain and at least one binding domain,
- 20 b) using at least one marker substrate which binds to the catalytic domain and is converted,
- c) using at least one substrate which can bind to the catalytic domain and to the binding domain,
- 25 d) incubating said protein, the marker substrate and the substrate with the substance or ligand, and
- e) determining whether the marker substrate is converted by the protein.

- 30 Suitable probes are, for example, enzymes which contain at least one catalytic domain and at least one binding domain. The enzymes may, however, also contain 1 to 8 catalytic domains and 1 to 8 binding domains, with the most

common enzymes containing 1, 2, 3 or 4 catalytic domains or binding domains. The substrate is a compound which can bind both to the binding domain and to the catalytic domain and which is chemically converted by the catalytic domain but not chemically modified by the binding domain. The marker substrate is a
5 compound which is chemically different from the substrate, is chemically converted by the catalytic domain and allows monitoring of the conversion reaction.

If a test substance binds essentially reversibly or irreversibly to the binding
10 domain of the protein, it may be regarded as an inhibitor of the binding domain of the protein. Not to be limited by theory, inhibition of the binding domain of the protein decreases competition for the catalytic domain of the protein between the marker substrate and the substrate—favoring conversion of the marker substrate—because the substrate generally binds to the binding domain before it is
15 transformed in the catalytic domain of the protein. A desired inhibitor or ligand, would not significantly prevent chemical conversion of the marker substrate by the catalytic domain of the protein because the marker substrate does not need to bind to the binding domain of the protein to be converted.

In summary, when an inhibitor or ligand is present, conversion of the substrate by
20 the protein decreases but no inhibition of marker substrate occurs. Conversely, when the test substance is not an effective inhibitor or ligand, there is competition between the marker substrate and the substrate for the catalytic domain of the protein with the end result that conversion of the marker substrate is reduced.

25 An example of a suitable protein is the enzyme collagenase which comprises at least two covalently linked enzyme domains, one which binds collagen and another which has proteolytic capabilities and cuts the collagen strand. This catalytic proteolytic domain, whether part of the full-length enzyme, i.e. the naturally occurring active enzyme, or just a proteolytic domain prepared by
30 recombination, has the ability to cleave marker substrates of different types.

Examples of suitable marker substrates for collagenase are peptides provided with fluorescent markers or UV markers. Examples of marker substrates for enzymes of this kind are (7-methoxycoumarin-4-yl)acetyl-Pro-Leu-Gly-Leu-(3-[2,4-dinitrophenyl]-L-2,3-diaminopropionyl)-Ala-Arg-NH₂, (C.G. Knight et al, FEBS Letters 296:263-266 (1992)), Dnp-Pro-β-cyclohexyl-Ala-Gly-Cys(Me)-His-Ala-Lys(N-Me-Abz)-NH₂ (D. M. Bickett et al., Analytical Biochemistry 212:58-64(1993)); Mca-Pro-Cha-Gly-Nva-His-Ala-Dpa-NH₂ (V. Knäuper et al., JBC 271/3, 1544-1550 (1996)); Mca-Arg-Pro-Lys-Pro-Val-Glu-Nva-Trp-Arg-Lys-Dnp-NH₂ (H. Nagase et al., JBC 269/33, 20952-20957 (1994)); Dnp-Arg-Pro-Lys-Pro-Leu-Ala-Nva-Trp-NH₂ (L. Niedzwiecki et al., Biochemistry 31:12618-12623 (1992)) (D. M. Bickett et al., Analytical Biochemistry 212:58-64 (1993); C.G. Knight, et al, FEBS Letters 296:263-266 (1992); V. Knäuper et al., JBC 271/3, 1544-1550 (1996); H. Nagase et al., JBC 269/33, 20952-20957 (1994); L. Niedzwiecki et al., Biochemistry 31:12618-12623 (1992)), or radiolabeled peptides.

Detection of marker substrates is not limited to fluorescence, UV/Vis, or radioactivity measurements but includes any suitable method known in the art for this purpose, such as HPLC, GC among others.

A suitable substrate for collagenase is collagen. Collagen is a proline-rich structural protein (scleroprotein) and the major component of mesenchymal intercellular supportive substances, which protects against enzymatic attacks. Three protein chains with left-handed helical structure are twisted into a right-handed triple helix (superhelix). 18 collagen types (collagen type I to type XVIII) have been identified, which can be classified according to structure or function into fibrillar, fibril-associated and nonfibrillar collagens.

A particularly suitable substrate for collagenase is type II collagen. This collagen supports the cartilage matrix in joints. In certain diseases such as osteoarthritis and rheumatism the joints are destroyed, in particular due to proteolytic degradation of collagen by collagenases. Inhibitors of enzymes of this kind are

known but have the disadvantage of attacking the catalytic domain of the enzyme (K. U. Weithmann et al., *Inflamm. Res.* 46:246-252 (1997)). Said catalytic domain is part, in a similar structure, of many enzymes and, as a result, the inhibitors act in an unwanted manner upon many enzymes, including those having a vital function (I. Massova et al., *The FASEB Journal* 12:1075-1095 (1998)). It is the object of the present invention to make better inhibitors, which have a higher specificity for the particular enzymes, available for medicine.

When a marker substrate and type II collagen were incubated with the enzyme collagenase and the marker substrate conversion was determined, it was surprisingly found that the marker substrate conversion was strongly inhibited. This inhibition could be overcome only by large amounts of marker substrate. However, said inhibition of marker substrate conversion did not occur on incubation of only the catalytic domain of collagenase with collagen II and marker substrate.

Surprisingly, it is possible in the method of the invention to identify substances which essentially abolish the inhibition of marker substrate conversion on incubation of complete collagenase with a marker substrate, type II collagen and the test substance. This effect does not appear when only the catalytic domain of collagenase is used instead of complete collagenase.

In contrast, previously reported enzyme inhibitors (K. U. Weithmann et al., *Inflamm. Res.* 46:246-252 (1997)) cause increased inhibition of marker substrate conversion catalyzed both by the complete enzyme and by the catalytic domain.

Examples of a "protein which contains at least one binding domain and at least one catalytic domain" are the members of the matrixins, the enzymes gelatinase, collagenase-1, neutrophil collagenase or matrix metalloproteinase of type 13 (I. Massova et al., *The FASEB Journal* 12, 1075-1095 (1998)).

09376091 "060801

5

10

15

- 20

25

5 fluorophenyl)-4-phenyl-3-butenyl]piperidine and/or of a stereoisomeric form of 4-(diphenyl-methylene)-1-[4-(p-fluorophenyl)-4-phenyl-3-butenyl]piperidine together with a pharmaceutically suitable and physiologically tolerated carrier, additive and/or other active ingredients and excipients.

10 In addition to the compound 4-(diphenylmethylene)-1-[4-(p-fluorophenyl)-4-phenyl-3-butenyl]piperidine, owing to the pharmacological properties, protamine sulfate and also oligosaccharides and tetrasaccharides, which are obtained from heparin degradation, are also suitable for the prophylaxis and therapy of all those disorders whose course involves increased activity of matrix-degrading enzymes
15 such as collagenases, metalloproteinases or aggrecanase. This includes degenerative joint disorders such as osteoarthroses, spondyloses, chondrolysis after joint trauma or prolonged immobilization of joints after meniscus or patella injuries or tearing of ligaments. This further includes also disorders of the connective tissue such as collagenoses, periodontal disorders, wound-healing
20 disturbances and chronic disorders of the locomotor apparatus such as inflammatory, immunologically or metabolically caused acute and chronic arthritis, arthropathies, myalgias and disorders of bone metabolism. Furthermore, 4-(diphenylmethylene)-1-[4-(p-fluorophenyl)-4-phenyl-3-butenyl]piperidine is suitable for the treatment of ulceration, atherosclerosis and stenoses. In addition,
25 4-(diphenylmethylene)-1-[4-(p-fluorophenyl)-4-phenyl-3-butenyl]piperidine is suitable for the treatment of inflammations, cancers, formation of tumor metastases, cachexia, anorexia and septic shock.

The pharmaceutical of the invention is generally administered orally or parenterally. Rectal or transdermal administration is also possible.

The invention also relates to a method for preparing a pharmaceutical, which comprises bringing 4-(diphenylmethylene)-1-[4-(p-fluorophenyl)-4-phenyl-3-

5 Examples of suitable solid or liquid pharmaceutical preparation forms are granules, powders, coated tablets, tablets, (micro)capsules, suppositories, syrups, elixirs, suspensions, emulsions, drops or injectable solutions and also products with protracted release of active ingredient, whose preparation makes use of common auxiliaries such as carriers, disintegrants, binders, coating agents, swelling agents, glidants or lubricants, flavorings, sweeteners and solubilizers. Frequently used excipients which may be mentioned are magnesium carbonate, titanium dioxide, lactose, mannitol and other sugars, talc, lactoprotein, gelatin, starch, cellulose and its derivatives, animal and vegetable oils such as cod liver oil, sunflower, peanut or sesame oil, polyethylene glycol and solvents such as, for example, sterile water and mono- or polyhydric alcohols such as glycerol.

compound of the invention, 4-(diphenylmethylene)-1-[4-(p-fluorophenyl)-4-phenyl-3-butenyl]piperidine. In the case of solid dosage units such as tablets, capsules, coated tablets or suppositories, said dose can be up to about 1000 mg, but is typically from about 50 to about 300 mg, and in the case of injection solutions in ampoule form up to about 300 mg, but typically from about 10 to about 100 mg. For the treatment of an adult patient about 70 kg in weight, daily doses of about 20 mg to about 1000 mg, typically from about 100 mg to about 500 mg, of active ingredient are indicated. Under certain circumstances, however, higher or lower daily doses may also be appropriate. The daily dose can be administered either by single administration in the form of an individual dosage unit or else of several smaller dosage units or by multiple administration of divided doses at particular intervals.

The invention further relates to an inhibitor which is obtainable through detection by the method of the invention.

TE0990" TE092860

Examples

Preparation of the COMPONENTS:

5 COMPONENT A)

Enzyme solution:

10 MMP-13 was obtained as a commercial product (Cat.No. HM 13110010) from
Invitex GmbH, Berlin, Germany, activated according to the manufacturer's
instruction, and diluted to 10 µg/10 ml with TCB buffer. TCB buffer was prepared
by dissolving 10 mM tris(hydroxymethyl)aminomethane (Sigma-Aldrich Chemie
15 GmbH, Deisenhofen, Germany) in water and adjusting the pH to 7.5 with HCl.
100 mM CaCl₂·2H₂O (Merck KGaA, Darmstadt, Germany) and 0.05% of Brij 35
solution, 30% (w/v) (Sigma-Aldrich Chemie GmbH, Deisenhofen, Deutschland),
are added to said solution.

COMPONENT B)

Marker substrate solution

20 A solution of 10 mmol/l (7-methoxycoumarin-4-yl)acetyl-Pro-Leu-Gly-Leu-(3-[2,4-
dinitrophenyl]-L-2,3-diaminopropionyl)-Ala-Arg-NH₃ (Bachem Biochemica GmbH,
Heidelberg, Germany) in dimethyl sulfoxide (DMSO) (Riedel-de Haen AG,
Seelze, Germany) was diluted 1 : 40 (v/v) with water.

COMPONENT C)

25 Collagen II solution

10 mg of human collagen II (Biocon, Potsdam, Germany) were dissolved in 2400
µl of 10mmol/l acetic acid (72 hours at 4° C), and then 1300 µl of a sodium
bicarbonate solution (250 mmol/l) CaCl₂·2H₂O (Merck KGaA, Darmstadt,
Germany) were added dropwise.

30

COMPONENT D)

Test substance (suspected inhibitor or ligand), typically dissolved in water at an appropriate concentration relative to the amount of enzyme used.

5 Test protocol

The components A) (25 μ l), B) (5 μ l), C) (10 μ l) and D) (10 μ l) were mixed in a total volume of 50 μ l, and fluorescence was measured in a commercial spectrofluorimeter after 15 minutes (excitation at 330 nm, emission at 390 nm,

10 Spectrafluor plus, Tecan Deutschland GmbH, Crailsheim, Germany).

15 Enzyme and inhibitor were preincubated at room temperature for 15 min. The reaction was started by adding 5 μ l of marker substrate solution (component B at 25 μ M).

The fluorescence in example 1 was set at 100%, and the measured values of the other examples are based on this value. Table 1 shows the results.

20 Table 1

	Example 1	Example 2	Example 3	Example 4	Example 5	Example 6
Component A	yes	yes	Yes	yes	yes	yes
Component B	yes	yes	Yes	yes	yes	yes
Component C	no	yes	Yes	yes	yes	yes
Component D	no	no	200 μ mol/l substance (I)	200 μ mol/l doxycycline	50 μ g/well tetra- saccharide	60 μ g/well protamine sulfate
Measured value	100%	49%	59%	0%	72%	73%

Examples 3, 5 and 6 illustrate the method of the invention, which makes it possible to detect novel inhibitors or ligands. These are distinguished by producing fluorescence values of 50-99% at a concentration of up to 500 $\mu\text{mol/l}$, typically 50 $\mu\text{mol/l}$ or less, in particular also starting from 1 nmol/l , and in exceptional cases
5 down to 0.1 nmol/l .

Example 1:

In general, inhibitor or ligands of the invention can be detected by comparing the transformation of the marker substrate in the presence of the test substance with
10 the corresponding transformation occurring in appropriate control mixtures. One such control mixture contained the enzyme and the substrate marker but no substrate or test substance (e.g., Example 1 in Table 1). The transformation of the marker substrate in this control mixture indicated the maximum conversion possible for the marker substrate because there was no inhibition of the catalytic
15 domain.

Example 2:

Another control mixture contained the enzyme the substrate marker and the substrate but no test substance (e.g., Example 2 in Table 1). Transformation of
20 the marker substrate in this second control mixture reflected competitive inhibition of the catalytic domain by both the substrate and the marker substrate.

If conversion of the marker substrate in the presence of a test substance is between the values obtained with these two control mixtures, an inhibitor or
25 ligand of the invention has been found.


Conversely, if transformation of the marker substrate is lower than the value obtained with the second control mixture, then the test substance is likely competing with both the substrate and the marker substrate for binding to the
30 catalytic site.

09376091 060301

In general, transformation of the marker substrate can be detected by a variety of methods known in the art that include, but are not limited to, the determination of the rate of transformation of the marker substrate at a given moment or the determination of the overall conversion of the marker substrate after incubation for a suitable period of time.

Example 3:

200µmol/l of the substance of the invention, 4-(diphenylmethylene)-1-[4-(p-fluorophenyl)-4-phenyl-3-butenyl]piperidine, (I)



produced a value of 59%.

Example 4

Those substances producing measured values of 49% or less, decreasing down to zero, have to be distinguished from the substances of the invention. For example, the prior art, 200 µmol/l doxycycline (K. U. Weithmann et al., Inflamm. Res. 46:246-252 (1997)) produced a value of 0.

Example 5

50µg/50µl tetrasaccharide GT8021 (Neoparin, Inc., 14274 Wicks Blvd, San Leandro, CA 94577). According to the invention, 50µg/50µl tetrasaccharide produced a value of 72%.

5

- 14 -